

Die provisorische und definitive Implantatversorgung ohne Abutmentwechsel im digitalen Workflow

# Zwei Scans, aber nur ein Abutment

Ein Beitrag von Dr. Robert Schneider MSc

## Literaturangabe

- [1] Beuer F, Schweiger J, Edelhoff D, Güth J, Stimmelmayer M. Das Münchener Implantatkonzept MIC - Eine praxisreife Kombination von Intraoralscanner und digitaler Fertigung. Dtsch Zahnärztl Z 2014;69:18-24.
- [2] Degidi M, Nardi D, Piattelli A: One abutment at one time: non-removal of an immediate abutment and its effect on bone healing around subcrestal tapered implants. Clin Oral Implants Res 2011;22:1303-1307.
- [3] Kern M, Lehmann F. Influence of surface conditioning on bonding to polyetheretherketon (PEEK). Dental materials : official publication of the Academy of Dental Materials 2012; 28: 1280-1283.
- [4] Koutouzis T, Richardson J, Lundgren T. Comparative soft and hard tissue responses to titanium and polymer healing abutments. The Journal of oral implantology 2011; 37 Spec No: 174-182.
- [5] Schmidlin PR, Stawarczyk B, Wieland M, Attin T, Hammerle CH, Fischer J. Effect of different surface pre-treatments and luting materials on shear bond strength to PEEK. Dental materials : official publication of the Academy of Dental Materials 2010; 26: 553-559.
- [6] Verri C, Vavassori V, Borgonovo A E, Re D, Maiorana C: Einsatz von Hochleistungspolymeren (BioHPP) zur Rehabilitation bei oralen Implantaten. Scientific & Clinical Cases Sofortige Einzelzahnversorgung 2015; 16-17, bredent medical ISBN 978-3-9817105-4-0.
- [7] Volpe S, Verrocchi D, Andersson P, Gottlow J, Sennerby L. Comparison of early bacterial colonization of PEEK and titanium healing abutments using real-time PCR. App Oss Res. 2008; 6: 54-62.

