

Auswahl von Materialien in der restaurativen Zahnheilkunde

# Entscheidungskriterien für die tägliche Praxis

Ein Beitrag von Prof. Dr. Gerwin Arnetzl



1. Acunzo R, Limioli E, Pagni G, Dudaite A, Consonni D, Rasperini G. Gingival Margin Stability After Mucogingival Plastic Surgery. The Effect of Manual Versus Powered Toothbrushing: A Randomized Clinical Trial. *J Periodontol* 2016; 87: 1186-1194.
2. Andlin-Sobocki A, Bodin L. Dimensional alterations of the gingiva related to changes of facial/lingual tooth position in permanent anterior teeth of children. A 2-year longitudinal study. *J Clin Periodontol* 1993; 20: 219-224.
3. Aroca S, Keglevich T, Nikolidakis D, Gera I, Nagy K, Azzi R, Etienne D. Treatment of class III multiple gingival recessions: a randomized-clinical trial. *J Clin Periodontol* 2010; 37: 88-97.
4. Baker DL, Seymour GJ. The possible pathogenesis of gingival recession. A histological study of induced recession in the rat. *J Clin Periodontol* 1976; 3: 208-219.
5. Bouchard P, Malet J, Borghetti A. Decision-making in aesthetics: root coverage revisited. *Periodontol* 2000 2001; 27: 97-120.
6. Burkhardt R, Hammerle CH, Lang NP, Research Group on Oral Soft Tissue B, Wound H. Self-reported pain perception of patients after mucosal graft harvesting in the palatal area. *J Clin Periodontol* 2015; 42: 281-287.
7. Burkhardt R, Lang NP. Coverage of localized gingival recessions: comparison of micro- and macrosurgical techniques. *J Clin Periodontol* 2005; 32: 287-293.
8. Cairo F, Cortellini P, Pilloni A, Nieri M, Cincinelli S, Amunni F, Pagavino G, Tonetti MS. Clinical efficacy of coronally advanced flap with or without connective tissue graft for the treatment of multiple adjacent gingival recessions in the aesthetic area: a randomized controlled clinical trial. *J Clin Periodontol* 2016; 43: 849-856.
9. Cairo F, Cortellini P, Tonetti M, Nieri M, Mervelt J, Cincinelli S, Pini-Prato G. Coronally advanced flap with and without connective tissue graft for the treatment of single maxillary gingival recession with loss of inter-dental attachment. A randomized controlled clinical trial. *J Clin Periodontol* 2012; 39: 760-768.
10. Cairo F, Nieri M, Pagliaro U. Efficacy of periodontal plastic surgery procedures in the treatment of localized facial gingival recessions. A systematic review. *J Clin Periodontol* 2014; 41 Suppl 15: S44-62.
11. Cairo F, Pagliaro U, Nieri M. Treatment of gingival recession with coronally advanced flap procedures: a systematic review. *J Clin Periodontol* 2008; 35: 136-162.
12. Chambrone L, Chambrone D, Pustigliani FE, Chambrone LA, Lima LA. Can subepithelial connective tissue grafts be considered the gold standard procedure in the treatment of Miller Class I and II recession-type defects? *J Dent* 2008; 36: 659-671.
13. Checchi L, Daprile G, Gatto MR, Pelliccioni GA. Gingival recession and toothbrushing in an Italian School of Dentistry: a pilot study. *J Clin Periodontol* 1999; 26: 276-280.
14. Coatoam GW, Behrents RG, Bissada NF. The width of keratinized gingiva during orthodontic treatment: its significance and impact on periodontal status. *J Periodontol* 1981; 52: 307-313.
15. Cortellini P, Pini Prato G. Coronally advanced flap and combination therapy for root coverage. Clinical strategies based on scientific evidence and clinical experience. *Periodontol* 2000 2012; 59: 158-184.
16. Dorfman HS, Kennedy JE, Bird WC. Longitudinal evaluation of free