

Von der Vitalerhaltung bis zur Pulpektomie

Endodontie im Milchgebiss

Ein Beitrag von Dr. Sven O. Pabel und Prof. Dr. Michael Hülsmann



Literaturangabe

- [1] Aeinehchi M, Dadvand S, Fayazi S, Bayat-Movahed S. Randomized controlled trial of mineral trioxide aggregate and formocresol for pulpotomy in primary molar teeth. *Int Endod J* 2007;40:261-267.
- [2] Ahmad IA, Pani SC. Accuracy of electronic apex locators in primary teeth: a meta-analysis. *Int Endod J* 2015;48:288-307.
- [3] Ahmed HM. Anatomical challenges, electronic working length determination and current developments in root canal preparation of primary molar teeth. *Int Endod J* 2013;46:1011-1022.
- [4] American Academy of Pediatric Dentistry (2014). Guideline on pulp therapy for primary and immature permanent teeth. *Reference Manual V36*;6:14-15.
- [5] Anthonappa RP, King NM, Martens LC. Is there sufficient evidence to support the long-term efficacy of mineral trioxide aggregate (MTA) for endodontic therapy in primary teeth? *Int Endod J* 2013;46:198-204.
- [6] Azar MR, Mokhtare M. Rotary Mtwo system versus manual K-file instruments: efficacy in preparing primary and permanent molar root canals. *Indian J Dent Res* 2011;22:363.
- [7] Barcelos R, Santos MP, Primo LG, LUIZ RR, Maia LC. ZOE paste pulpectomies outcome in primary teeth. *J Clin Ped Dent* 2011;42:241-248.
- [8] Beltrame AP, Triches TC, Sartori N, Bolan M. Electronic determination of root canal working length in primary molar teeth: an in vivo and ex vivo study. *Int Endod J* 2011;44:402-406.
- [9] Bjørndal L, Thylstrup A. A practicebased study on stepwise excavation of deep carious lesions in permanent teeth: a 1-year follow-up study. *Community Dent Oral Epidemiol* 1998;26: 122-128.
- [10] Botton G, Pires CW, Cadona FC, Machado AK, Azzolin VF, Cruz IB, Sagrillo MR, Praetzel JR. Toxicity of irrigating solutions and pharmacological associations used in pulpectomy of primary teeth. *Int Endod J* (epub ahead of print, doi:10.1111/iej.12509)
- [11] Burnett S, Walker J. Comparison of ferric sulfate, formocresol, and a combination of ferric sulfate/formocresol in primary tooth vital pulpotomies: A retrospective radiographic survey. *J Dent Child* 2002;69:44-48.
- [12] Cortes O, Garcia C, Perez L. Comparison between rotary and manual instrumentation in primary teeth. *J Clin Pediatr Dent* 2008;32:295-298.
- [13] Da Silva LA, Leonardo MR, Nelson-Filho P, Tanomaru JM. Comparison of rotary and manual instrumentation techniques on cleaning capacity and instrumentation time in deciduous molars. *J Dent Child* 2004;71:45-47.
- [14] De Paula VA, de Carvalho Ferreira D, Cavalcante FS, do Carmo FL, Rosado AS, Primo LG, dos Santos KR. Clinical signs and bacterial communities of deciduous necrotic root canals detected by PCR-DGGE analysis: research association. *Arch Oral Biol* 2014;59:848-854.
- [15] Erdem AP, Guven Y, Balli B, Ilhan B, Sepet E, Ulukapi I, Aktoren Q. Success rates of mineral trioxide aggregate, ferric sulfate, and formocresol pulpotomies: a 24-month study. *Pediatr Dent* 2011; 33:165-170.
- [16] Fuks A, Eidelman E, Pauker N. Root fillings with Endoflas in primary teeth: a retrospective study. *J Clin Pediatr Dent* 2002;27:41-46.
- [17] Fuks AB. Vital pulp therapy with new materials for primary teeth: new directions and treatment perspectives. *J Endod* 2008;34(Suppl):S18-24.
- [18] Grover R, Mehra M, Pandit IK, Srivastava N, Gugnani N, Gupta M. Clinical efficacy of various root canal obturating methods in primary teeth: a comparative study. *Eur J Paediatr Dent* 2013;14:104-108.
- [19] Heinrich-Weltzien R, Hickel R, Kühnisch J. Die Caries-profunda-Therapie im Milchgebiss. *Endodontie* 2011;20:139-148.
- [20] Holan G. Long-term effect of different treatment modalities for traumatized primary incisors presenting dark coronal discoloration with no other signs of injury. *Dent Traumatol* 2006;22:14-17.
- [21] Hori A, Poureslami HR, Parirokh M, Mirzazadeh A,

- Abbott P. The ability of pulp sensibility tests to evaluate the pulp status in primary teeth. *Int J Paediatr Dent* 2011; 21:441-445.
- [22] Hülsmann M, Wiegand A. Endodontie im Milch- und Wechselgebiss. In: Einwag J, Pieper K (Hrsg.). *Kinderzahnheilkunde*. 3. Aufl., Urban & Fischer. München-Jena 2008, 249-268.
- [23] Hülsmann M. Endometrie im Milchgebiss. *Endodontie* 2011;20:161-166.
- [24] Huth KC, Hajek-Al-Khatir N, Wolf P, Ilie N, Hickel R, Paschos E. Long-term effectiveness of four pulpotomy techniques: 3-year randomized controlled trial. *Clin Oral Invest* 2012;16:1243-1250.
- [25] Innes NPT, Evans DJP, Stirrups DR. Sealing caries in primary molars; randomized control trial, 5-year results. *J Dent Res* 2011;90:1405-1410.
- [26] Innes NPT, Stirrups DR, Evans DJP, Hall N. A novel technique using preformed metal crowns for managing carious primary molars in general practice – a retrospective analysis. *Brit Dent J* 2006; 200:451-454.
- [27] Ito IY, Matoba F, Paula-Silva FW, Bezerra LA, Leonardo MR, Nelson-Filho P. Microbial culture and checkerboard DNA-DNA hybridization assessment of bacteria in root canals of primary teeth pre- and post-endodontic therapy with a calcium hydroxide/chlorhexidine paste. *Int J Paediatr Dent* 2011;21: 353-360.
- [28] Katge F, Patil D, Poojari M, Pimpale J, Shitoot A, Rusaawat B. Comparison of instrumentation time and cleaning efficacy of manual instrumentation, rotary systems and reciprocating systems in primary teeth: an in vitro study. *J Indian Soc Pedod Prev Dent* 2014;32:311-316.
- [29] Katz A, Mass E, Kaufman AY. Electronic apex locator: a useful tool for root canal treatment in the primary dentition. *ASDC J Dent Child* 1996;63:414-417.
- [30] Klein U, Kleier DJ. Sodium hypochlorite accident in a pediatric patient. *Pediatr Dent* 2013;35:534-538.
- [31] Koshy S, Love RM. Endodontic treatment in the primary dentition. *Aust Endod J* 2004;30:59-68.
- [32] Kühnisch J, Heinrich-Weltzien R, Schäfer E. Endodontie im Milchgebiss. Wissenschaftliche Mitteilung der Deutschen Gesellschaft für Kinderzahnheilkunde (DGKiZ) und der Deutschen Gesellschaft für Zahnerhaltung (DGZ). *Endodontie* 2011a;20:167-172.
- [33] Kühnisch J, Hickel R, Heinrich-Weltzien R. Direkte Überkappung, Pulpotomie und Pulpektomie am Milchzahn. Indikationen, Kontraindikationen und klinischer Behandlungsablauf. *Endodontie* 2011b;20:149-160.
- [34] Kummer TR, Calvo MC, Cordeiro MM, de Souza Vieira R, Rocha MJ. Ex vivo study of manual and rotary instrumentation techniques in human primary teeth. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;105:e84-02.
- [35] Lange J, Bethke S, Stoll R, Pieper K, Jablonski-Momeni A. Der Einfluss verschiedener Parameter auf das Überleben von endodontischen Maßnahmen an Milchzähnen. *Oralprophylaxe & Kinderzahnheilkunde* 2012;34:20-25.
- [36] Lima CC, Conde Junnior AM, Moura RD, Moura MS, Lima MD, Moura LF. Biocompatibility of root filling pastes used in primary teeth. *Int Endod J* 2015;48:405-416.
- [37] Lima RA, Carvalho CB, Ribeiro TR, Fonteles CS. Antimicrobial efficacy of chlorhexidine and calcium hydroxide/camphorated paramonochlorophenol on infected primary molars: a split-mouth randomized clinical trial. *Quintessence Int* 2013;44:113-122.
- [38] Louwakul P, Prucksathamrongkul W. The effect of 2% chlorhexidine as root canal irrigant in pulpectomies of primary molars. *Pediatr Dent* 2012;34:e192-196.
- [39] Lucas Leite AC, Rosenblatt A, da Silva Calixto M, da Silva CM, Santos N. Genotoxic effect of formocresol pulp therapy of deciduous teeth. *Mutat Res* 2012;747:93-97.
- [40] Lustig JP, Schwartz-Arad D, Shapira A. Odontogenic cysts related to pulpotomized deciduous molars. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1999;87:499-503.
- [41] Magnusson BO, Sundell SO: Stepwise excavation of deep carious lesions in primary molars. *J Int Assoc Dent Child* 1977;8:36-40
- [42] Mahoney P: Two-dimensional patterns of human enamel thickness on deciduous (dm1, dm2) and permanent first (M1) mandibular molars. *Arch Oral Biol* 2010;55:115-126.
- [43] Makarem A, Ravandeh N, Ebrahimi M. Radiographic assessment and chair time of rotary instruments in the pulpectomy of primary second molar teeth: a randomized controlled clinical trial. *J Dent Res Dent Clin Dent Prospects* 2014;8:84-89.
- [44] Maroto M, Barberia E, Planells P, Garcia-Godoy F. Dentin bridge formation after mineral trioxide aggregate (MTA) pulpotomies in primary teeth. *Am J Dent* 2005;18:151-154.
- [45] Mörchen-Trinkle S, Biggel A. Endodontische Behandlung von Milchzähnen bei Aplasie bleibender Zähne. *Endodontie* 2014;23:179-187.
- [46] Moskovitz M, Tickotsky N, Ashkar H, Holan G. Degree of root resorption after root canal treatment with iodoform-containing filling material in primary molars. *Quintessence Int* 2012;43:361-368.
- [47] Nagaratna PJ, Shashikiran ND, Subbareddy VV. In vitro comparison of NiTi rotary instruments and stainless steel hand instruments in root canal preparations of primary and permanent molar. *J Indian Soc Pedod Prev Dent* 2006;24:186-191.
- [48] Ng FK, Messer LB. Mineral trioxide aggregate as a pulpotomy medicament: an evidence-based assessment. *Eur Arch Paediatr Dent* 2008;9:58-73.
- [49] Ochoa-Romero T, Mendez-Gonzalez V, Flores-Reyes

- H, Pozos-Guillen AJ. Comparison between rotary and manual techniques on duration of instrumentation and obturation times in primary teeth. *J Clin Pediatr Dent* 2011;35:359-363.
- [50] Oznurhan F, Unal M, Kapdan A, Ozturk C, Aksoy S. Clinical evaluation of apex locator and radiography in primary teeth. *Int J Paediatr Dent* 2015;25:199-203
- [51] Parirokh M, Torabinejad M. Mineral trioxide aggregate: a comprehensive literature review. Part III: clinical applications, drawbacks, and mechanism of action *J Endod* 2011;36:400-413,
- [52] Peng L, Ye L, Tan H, Zhou X. Evaluation of the formocresol versus mineral trioxide aggregate primary molar pulpotomy: a meta-analysis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102:e40-44.
- [53] Pinheiro SL, Araujo G, Bincelli I, Cunha R, Bueno C. Evaluation of cleaning capacity and instrumentation time of manual, hybrid and rotary instrumentation techniques in primary molars. *Int Endod J* 2012;45:379-385.
- [54] Pinto DN, de Sousa DL, Araújo RB, Moreira-Neto JJ. Eighteen-month clinical and radiographic evaluation of two root canal-filling materials in primary teeth with pulp necrosis secondary to trauma. *Dent Traumatol* 2011;27:221-224.
- [55] Pires CW, Botton G, Cadona FC, Machado AK, Azzolin VF, Cruz IB, Sagrillo MR, Praetzel JR. Induction of cytotoxicity, oxidative stress, and genotoxicity by root filling pastes used in primary teeth. *Int Endod J* (epub ahead of print; doi:10.1111/iej.12502).
- [56] Pitoni CM, Figueiredo MC, Araujo FB, Souza MA. Ethylenediaminetetraacetic acid and citric acid solutions for smear layer removal in primary tooth root canals. *J Dent Child* 2011;78:131-137.
- [57] Pramila R, Muthu MS, Deepa G, Farzan JM, Rodrigues SJ. Pulpectomies in primary mandibular molars: a comparison of outcomes using three root filling materials. *Int Endod J* 2015; e-pub ahead of print doi: 10.1111/iej.12478.
- [58] Ranly DM, Garcia-Godoy F. Current and potential pulp therapies for primary and young permanent teeth. *J Dent* 2000;28:153-161.
- [59] Ruiz-Esparza CL, Garrocho-Rangel, A, Gonzales Amaro AM, Florez-Reyes H, Pozos-Guillen AJ. Reduction in bacterial loading using 2% chlorhexidine gluconate as an irrigant in pulpectomized primary teeth: a preliminary report. *J Clin Pediatr Dent* 2011;35:265-270.
- [60] Schwendicke F, Paris S. Kariesexkavation: ein systematischer Überblick. *Dtsch Zahnärztl Z* 2014;69:456-466.
- [61] Seraj B, Ramezani G, Ghadimi S, Mosharrafian SH, Motahhary P, Safari M. In-vitro comparison of instrumentation time and cleaning capacity between endodontic handpiece and manual preparation techniques in primary molar teeth. *Minerva Stomatol* 2013;62:17-22.
- [62] Shirvani A, Asgary S. Mineral trioxide aggregate versus formocresol pulpotomy: a systematic review and meta-analysis of randomized clinical trials. *Clin Oral Invest* 2014;18:1023-1030.
- [63] Smail-Faugeron V, Courson F, Durieux P, Muller-Bolla M, Glenny AM, Fron Chabouis H. Pulp treatment for extensive decay in primary teeth. *Cochrane Database Syst Rev* 2014; doi: 10.1002/14651858.
- [64] Stephan B, Bimler M. Pulpotomie und Hemisektion von Milchmolaren zur Unterstützung des kieferorthopädischen Lückenschlusses bei Nichtanlage des zweiten unteren Prämolaren. *Endodontie* 2014;23:189-194.
- [65] Subramaniam P, Tabrez TA, Babu KL. Microbiological assessment of root canals following use of rotary and manual instruments in primary molars. *J Clin Pediatr Dent* 2013;38:123-127.
- [66] Tannure PN, Azevedo CP, Barcelos R, Gleiser R, Primo LG. Long-term outcomes of primary tooth pulpectomy with and without smear layer removal: a randomized split-mouth clinical trial. *Pediatr Dent* 2011;33:316-20
- [67] Tannure PN, Barcelos R, Portela MB, Gleiser R, Primo LG. Histopathologic and SEM analysis of primary teeth with pulpectomy failure. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;108:e29-e33.
- [68] Triches TC, de Figueiredo LC, Feres M, de Freitas SF, Zimmermann GS, Cordeiro MM. Microbial profile of root canals of primary teeth with pulp necrosis and periradicular lesion. *J Dent Child* 2014;81:14-19.
- [69] Tulsani SG, Chikkanarasaiah N, Bethur S. An in vivo comparison of antimicrobial efficacy of sodium hypochlorite and Biopure MTAD against enterococcus faecalis in primary teeth: a qPCR study. *J Clin Pediatr Dent* 2014 39:30-34.
- [70] UK National Clinical Guidelines in Paediatric Dentistry. *Int J Paediatr Dent* 2008;18 Suppl 1:20-8. doi: 10.1111/j.1365-263X.2008.00935.x.
- [71] Wiegand A, Hülsmann M. Pulpotomie im Milchgebiss – Eine Standortbestimmung. *Endodontie* 2005;14:177-192.
- [72] Wrbas KT, Kielbassa AM, Hellwig E. Microscopic studies of accessory canals in primary molar furcations. *ASDC J Dent Child* 1997;64:118-122.
- [73] Ximenes M, Triches TC, Beltrame AP, Hilgert LA, Cardoso M. Effect of endodontic irrigation with 1% sodium hypochlorite and 17% EDTA on primary teeth: a scanning electron microscope analysis. *Gen Dent* 2013;61:24-27.