

Five-year follow-up of immediate restorations with ceramic-reinforced PEEK

# A safe and predictable long-term treatment protocol

PROFESSOR JOSÉ EDUARDO MATÉ SÁNCHEZ DE VAL, MURCIA, SPAIN

## Literature

1. Araújo MG, Sukekava F, Wennström JL, Lindhe J. Tissue-modeling following implant placement in fresh extraction sockets. *Clin Oral Implants Res* 2006;17:615–624.
2. Berglundh T, Lindhe J. Dimension of the periimplant mucosa. Biological width revisited. *J Clin Periodontol* 1996;23:971–973.
3. Abdullah, M.R., Goharian, A., Abdul Kadir, M.R. & Wahit, M.U. (2015) Biomechanical and bioactivity concepts of polyetheretherketone composites for use in orthopedic implants-a review. *Journal of Biomedical Materials Research, Part A*. 103: 3689-3702.
4. Hermann JS, Buser D, Schenk RK, Schoolfield JD, Cochran DL. Biologic width around one- and two-piece titanium implants. *Clin Oral Implants Res* 2001;12:559–571.
5. Lazzara RJ, Porter SS. Platformswitching. A new concept in implant dentistry for controlling post-restorative bone levels. *Int J Periodontics Restorative Dent* 2006;26:9–17.
6. Gardner DM. Platform switching as a means to achieving implant esthetics. A case study. *NY State Dent J* 2005;71:34–37.