

Versorgung einer unilateralen Nichtanlage

Der kieferorthopädische Lückenschluss

Ein Beitrag von Dr. Susanna Isabel Richter, Dr. Eva-Maria Niehoff, Dr. Elisabeth Klang und Dr. Frauke Beyling



Literaturangabe

- [1] Al-Qabandi SM: Slot dimension of completely customized lingual orthodontic brackets – a comparative analysis. Master's Thesis Lingual Orthodontics. Hannover Medical School, Department of Orthodontics; 2013
- [2] Balmaceda D: Dimensionsgenaue Herstellung eines Finishingbogens in der Lingualtechnik. Master's Thesis Lingual Orthodontics. Hannover Medical School, Department of Orthodontics; 2018
- [3] Bauss O, Sadat-Khonsari R, Engelke W, Kahl-Nieke B: results of transplanting developing third molars as part of orthodontic space management. Part 1: clinical and radiographic results. *J Orofac Orthop* 2002; 63, 483-492
- [4] Bennett JC, McLaughlin RP: Kieferorthopädisches Management im Bereich des zweiten Unterkiefer-Prämolar. *J Orofac Orthop* 2001; 33:265-274
- [5] Bieber C: Bestimmung der Torquekapazität einer vollständig individuellen lingualen Apparatur neuester Generation. Master's Thesis lingual Orthodontics. Hannover Medical School, Department of orthodontics, 2013
- [6] Bjerklin K, Bennett J. The long-term survival of lower second primary molars in subjects with agenesis of the premolars. *Eur J Orthod* 2000; 22: 145-157
- [7] Bock N, Ruf S: Post-treatment occlusal changes in class II devision 2 subjects treated with the Herbst appliance. *Eur J Orthod* 2008, 30: 606-613
- [8] Bock N, Ruf S, Wiechmann D, Jilek T: Herbst plus lingual versus Herbst plus labial: a comparison of occlusal outcome and gingival health. *Eur J Orthod* 2016, 38: 478-484
- [9] Cash AC, Good SA, Curtis RV, McDonald F: An evaluation of slot size in orthodontic brackets - are standard as expected? *Angle Orthod* 2004, 74: 450-453
- [10] Cattaneo PM, Salih RA, Melson B: Labio-lingual root control of lower anterior teeth and canines obtained by active and passive self-ligating brackets. *Angle Orthod* 2013, 83: 691-697
- [11] Cronin RJ, Oesterle LJ, Ranly DM: Mandibular implants and the growing patient. *Int J Oral Maxillofac Implants* 1994, 9: 55-62
- [12] Czochrowska EM, Stenvik A, Bjerke B, Zachrisson B: Outcome of tooth transplantation: survival and success rates 17-41 years posttreatment. *Am J Orthod Dentofacial Orthop* 2002; 121: 110-119
- [13] Demling A, Dittmer MP, Schwestka-Polly R: Comparative analysis of slot dimension in lingual bracketr systems. *Head Face Med* 2009; 5: 27
- [14] Fanning EA: Third molar emergence in Bostonians. *Am J Phys Anthropol* 1962, 20: 339
- [15] Fines CD, Rebellato J, Saier M: Congenitally missing mandibular second premolars: treatment outcome with orthodontic space closure. *Am J Orthod Dentofacial Orthop* 2003; 126: 676-682
- [16] Fiorentino G, Melson B: Asymmtric mandibular space closure. *J Clin Orthod* 1996; 30: 519-523
- [17] Fischer-Brandis H, Orthuber W, Es-Souni M, Meyer S: Torque transmission between square ire and brackets as a function of measurement, form and hardness parameters. *J Orofac Orthop* 2000, 61: 258-265
- [18] Franchi L, Baccetti T, McNamara JAJ: Treatment and post-treatment effects of acrylic splint Herbst appliance therapy. *Am J Orthod Dentofacial Orthop* 1999; 115: 429-438
- [19] Gianelly A, Goldman H. Philadelphia: Lea and Febiger; 1971. *Biologic Basis of Orthodontics*.
- [20] Glassmann SD: Autogenous tooth transplantation in the treatment of malocclusion. *Dent Clin North Am* 1981; 25: 109-116
- [21] Grauer D, Proffit WR: Accuracy in tooth positioning with a fully customized lingual orthodontic appliance. *Am J Orthod Dentofacial Orthop* 2011, 140: 433-443
- [22] Haralabakis H: Observations on the time of eruption, congenital absence, and impaction of third molar teeth. *Trans Eur Orthod Soc* 1957, 33: 308-309
- [23] Hellman M: Our third molar teeth, their eruption, presence and absence. *Dent Cosmos* 1936; 78: 750-762
- [24] Hinz R: Behandlungsmisserfolge durch unterlassene Ausgleichsextraktionen. *J Orofac Orthop* 1988; 49: 463-465
- [25] Ith-Hansen K, Kjaer I: Persistence of deciduous molars in subjects with agenesis of the second premolars. *Eur J Orthod* 2000; 22: 239-243
- [26] Joondeph DR, McNeill W: Congenitally absent second premolars: An interceptive approach. *Am J Orthod Dentofacial Orthop* 1971; 59: 50-66
- [27] Kalwizki M, Ney T, Göz G: Transplantation of a lower bicuspid after traumatic loss of three upper incisors. *J Orofac Orthop* 2003; 64: 57-66
- [28] Katzorke M: Okklusale Stabilität und Einstellung von Weisheitszähnen nach Lückenschluss im Unterkiefer von distal mit einer vollständig individuellen lingualen und einer Herbst-Apparatur. Master's Thesis lingual Orthodontics. Hannover Medical School, Department of orthodontics, 2018

- [29] Kim YH, Yang SM, Kim S, Lee JY Kim KE, Gianelly AA, Kyung SH: Midpalatal miniscrews for orthodontic anchorage: factors affecting clinical success. *Am J Orthod Dentofacial Orthop.* 2010; 137: 66-72
- [30] Kokich VO: Congenitally missing teeth: orthodontic management in the adolescent patient. *Am J Orthod Dentofacial Orthop.* 2002; 121: 594-595
- [31] Kurol J, Thilander B: Infraocclusion of primary molars with aplasia of the permanent successors. *Ang Orthod.* 1984; 54: 283-294
- [32] Kusy R, Whitley J: Assessment of second-order clearances between orthodontic archwires and bracket slots via the critical contact angle for binding. *Angle Orthod.* 1999; 69: 71-80
- [33] Lekholm U: The use of osseointegrated implants in growing jaws. *Int J Oral Maxillofac Implants.* 1993; 8: 243-244
- [34] Maeda A, Sakoguchi Y, Miyawaki S: Patient with oligodontia treated with a miniscrew for unilateral mesial movement of the maxillary molars and alignment of an impacted third molar. *Am J Orthod Dentofacial Orthop.* 2013; 144: 430-440
- [35] Mamopoulos A, Hägg U, Schröder U, Hansen K: Agenesis of mandibular second premolars. Spontaneous space closure after extraction therapy: a 4-year follow-up. *Eur J Orthod.* 1996; 18: 589-600
- [36] Meling TR, Odegaard J, Segner D: On bracket slot height: A methodologic study. *Am J Orthod Dentofacial Orthop.* 1998; 113: 387-393
- [37] Metzner R, Schwestka-Polly R, Helms HJ, Wiechmann D: Comparison of anchorage reinforcement with temporary anchorage devices or a Herbst appliance during lingual orthodontic protraction of mandibular molars without maxillary counterbalance extraction. *Head Face Med.* 2015; 11: 22
- [38] Neugebauer J, Stellzig A, Zöller JE: Implantologie und Kieferorthopädie: Synergie oder Alternative bei der Therapie von Nischenanlagen? *Stomatol.* 1999; 96:65-70
- [39] Odman J, Gröndahl K, Lekholm U, Thilander B: the effect of osseointegrated implants on the dento-alveolar development. A clinical and radiographic study in growing pigs. *Eur J Orthod.* 1991; 13: 279-286
- [40] Oesterle LJ, Cronin RJ, Ranly DM: Maxillary implants and the growing patients. *Int J Oral Maxillofac Implants.* 1993; 8: 377-387
- [41] Pancherz H, Hansen K: Mandibular anchorage in Herbst treatment. *Eur J Orthod.* 1988; 10: 149-164
- [42] Pauls A: Therapeutic accuracy of the completely customized lingual appliance WIN. Master's thesis Lingual Orthodontics. Hannover Medical School, Department of Orthodontics; 2015
- [43] Pauls AH: Therapeutic accuracy of individualized brackets in lingual orthodontics. *J Orofac Orthop.* 2010; 71: 348-361
- [44] Paulsen HU, Andreasen JO: Eruption of premolars subsequent to autotransplantation. A longitudinal radiographic study. *Eur J Orthod.* 1998; 20: 45-55
- [45] Postema RS: The clinical management of congenitally absent and malformed maxillary lateral incisors and second bicuspid. *J Gen Orthod.* 1994; 5: 14-19
- [46] Schopf P: Aplasie. In: Schopf P, Hrsg. *Curriculum der Kieferorthopädie.* Berlin: Quintessenz, 1991: 416-424
- [47] Thalheim A, Schwestka-Polly R: Clinical realisation of a setup in lingual orthodontics. *Inf Orthod Kieferorthop.* 2008; 40: 277-282
- [48] Thilander B, Odman J, Lekholm U: Orthodontic aspects of the use of oral implants in adolescents: a 10-year follow-up study. *Eur J Orthod.* 2001; 23: 715-731
- [49] Weschler D, Pancherz H: Efficiency of three mandibular anchorage forms in Herbst treatment: a cephalometric investigation. *Angle Orthod.* 2005; 75: 23-27
- [50] Wheeler RC., Wheeler's atlas of tooth form, 1984 Philadelphia WB Saunders Co.
- [51] Wiechmann D: A new bracket system for lingual orthodontic treatment. Part 1: theoretical background and development. *J Orofac Orthop.* 2002; 63: 234-245
- [52] Wiechmann D, Schwestka-Polly R, Pancherz H, Hohoff A: Control of mandibular incisors with the combined Herbst and completely customized lingual appliance- a pilot study. *Head Face Med.* 2010; 6:3
- [53] Wolf M: Posttherapeutische Veränderungen unter permanenter Retention- Bietet die Verwendung von vollständig individuellen lingualen Apparaturen Vorteile? Master's Thesis Lingual Orthodontics. Hannover Medical School, Department of Orthodontics; 2017
- [54] Van der Veen MH, Attin R, Schwestka-Polly R, Wiechmann D: Caries outcomes after orthodontic treatment with fixed appliances: do lingual brackets make a difference? *Eur J Oral Sci.* 2010; 118: 298-303
- [55] Van Laeken R, Martin CA, Dischinger T, Razmus T, Ngan P: treatment effects of the edgewise Herbst appliance: A cephalometric and tomographic investigation. *Am J Orthod Dentofacial Orthop.* 2006; 130: 582-593
- [56] Vu J, Pancherz H, Schwestka-Polly R, Wiechmann D: Correction of class II, Division 2 malocclusion using a completely customized lingual appliance and the Herbst device. *J Orofac Orthop.* 2012; 73: 225-235
- [57] Zimmer B, Schelper I, Seifi-Shirvandeh N: Localized orthodontic space closure for unilateral aplasia of lower second premolars. *Eur J Orthod.* 2007; 29: 210-216