


Management und Heilung periapikaler Läsionen

Zahnerhalt nach endodontischer Therapie

Ein Beitrag von PD Dr. Ralf Krug, Würzburg, und Dr. Sebastian Reich, Eichstätt



Literaturangabe

1. McCabe PS, Dummer PM. Pulp canal obliteration: an endodontic diagnosis and treatment challenge. *Int Endod J* 2012;45:177-197.
2. Krasner P, Rankow HJ. Anatomy of the pulp-chamber floor. *J Endod* 2004;30:5-16.
3. Hülsmann M, Schäfer E. Probleme der Desinfektion. In: Hülsmann M, Schäfer E (Hrsg.) „Probleme in der Endodontie. Prävention, Identifikation und Management“. Quintessenz, Berlin, 2. Auflage 2019:255-289.
4. Baugh D, Wallace J. The role of apical instrumentation in root canal treatment: a review of the literature. *J Endod* 2005;31:333-340.
5. Bystrom A, Sundqvist G. Bacteriologic evaluation of the efficacy of mechanical root canal instrumentation in endodontic therapy. *Scand J Dent Res* 1981;89:321-328.
6. Orstavik D. Time-course and risk analyses of the development and healing of chronic apical periodontitis in man. *Int Endod J* 1996;29:150-155.
7. Kruse C, Spin-Neto R, Wenzel A, Kirkevang LL. Cone beam computed tomography and periapical lesions: a systematic review analysing studies on diagnostic efficacy by a hierarchical model. *Int Endod J* 2015;48:815-828.
8. de Paula-Silva FW, Wu MK, Leonardo MR, da Silva LA, Wesselink PR. Accuracy of periapical radiography and cone-beam computed tomography scans in diagnosing apical periodontitis using histopathological findings as a gold standard. *J Endod* 2009;35:1009-1012.